

## Original Article

## HEPATOPROTECTIVE AND ANTIOXIDATIVE EFFECTS OF ALLIUM SATIVUM VAR CHINESE EXOTIC ON ACETAMINOPHEN INDUCED ACUTE HEPATITIS IN MALE ALBINO RATS

Sauda Usmani, Aysha Zaheer and Hamid Javaid Qureshi

**Objective:** To determine the hepatoprotective and antioxidative effects of ethanolic extract of *Allium sativum* var Chinese exotic on acetaminophen induced hepatotoxicity in male albino rats.

**Methods:** This study was carried out on 90 male albino rats. A single intraperitoneal dose of acetaminophen 750mg/kg was used to induce oxidative stress and hepatotoxicity. The rats were randomly divided into three groups of thirty each. Group A was given normal saline (negative control); group B was administered hepatotoxic dose of acetaminophen (positive control); group C (experimental) was pretreated with *Allium sativum* var Chinese exotic extract for 7 days before receiving hepatotoxic dose of acetaminophen (Experimental). Serum ALT, AST, ALP, total proteins, albumin and glutathione peroxidase levels in each group were estimated from terminal blood sampling done 24 hours after acetaminophen administration under ether anesthesia.

**Results:** *Allium sativum* var Chinese exotic manifested hepatoprotective and antioxidative effects by producing highly significant ( $p=0.000$ ) reduction in serum ALT and AST, but no significant ( $p=0.335$ ) reduction in serum ALP levels. This garlic extract also produced highly significant ( $p=0.000$ ) increase in serum albumin and significant increase in serum total proteins ( $p=0.027$ ) and glutathione peroxidase ( $p=0.025$ ) levels.

**Conclusions:** *Allium sativum* var Chinese exotic has potent hepatoprotective and antioxidative potential.

**Keywords:** *Allium sativum*, Chinese exotic, glutathione peroxidase, antioxidative, hepatoprotective, acetaminophen.


### Introduction

Acute hepatitis results in massive necrosis of liver cells leading to severe impairment of liver functions.<sup>1</sup> An estimated 1600 cases of acute hepatic failure occur each year in United States. Acetaminophen toxicity is the most common cause, accounting for at least 45% of the cases.<sup>2</sup>

Acetaminophen (AAP), which is also named paracetamol, is a commonly used antipyretic and analgesic. Overdose of acetaminophen can lead to acute liver injury and histopathological changes characterized by centrilobular necrosis.<sup>3</sup> Chronic alcohol use may greatly increase susceptibility to hepatotoxicity from acetaminophen because of depleted glutathione stores.<sup>4</sup>

Treatment of healthy adults with acetaminophen taken at the maximum daily recommended dose of 4g for 4 or more days frequently cause elevations in serum aminotransferases which often persist when acetaminophen concentrations are no longer measurable in plasma.<sup>5</sup> In some patients, chronic ingestion of therapeutic doses may produce hepatic necrosis and hepatitis, which persist long after the drug has been discontinued.<sup>6</sup> In most cases, there is no effective treatment other than stopping

the drug and providing general supportive care. N-acetylcysteine (NAC) has been used, as an antidote for acetaminophen toxicity.<sup>7</sup> The oxidative metabolite of acetaminophen is more toxic than the drug. Hepatotoxic doses of paracetamol deplete the normal levels of hepatic glutathione. The hepatic cytochrome P450 enzyme system metabolizes paracetamol, forming NAPQI (*N*-acetyl-*p*-benzoquinone imine). NAPQI is then irreversibly conjugated with the sulfhydryl groups of glutathione. Conjugation depletes glutathione, a natural antioxidant. The highly reactive active metabolite NAPQI appears to mediate much of the acetaminophen-related damage to liver tissue by forming covalent bonds with cellular proteins and subsequent activation of inflammatory mediator TNF- $\alpha$  that in turn contribute to tissue necrosis.<sup>8</sup> (Fig-1) *Allium sativum*, or "garlic" is widely used in culinary preparations.<sup>9</sup> Two varieties of *Allium sativum* grown in Punjab are Chinese (exotic), *Lehsan Gulabi* (local).<sup>10</sup> Traditional uses of *Allium sativum* include; use in intestinal disorders, diarrhea, flatulence, worms, respiratory infections, skin diseases, wounds, symptoms of aging, headache, flu, sore throat, fever and otitis media.<sup>5</sup>



Garlic contains sulfur-containing constituents like  $\gamma$ -glutamyl-S-alkyl-L-cysteine and S-alkyl-L-cysteine, sulfoxides, allicin, steroidal glycosides, lectins, prostaglandins, fructan, pectin, essential oil, adenosine, vitamins B<sub>1</sub>, B<sub>2</sub>, B<sub>6</sub>, C and E, biotin, nicotinic acid, fatty acids, glycolipids, phospholipids, anthocyanins, flavonoids, phenolics and essential amino acids. Allicin and other thiosulfates instantly decompose to other compounds, such as diallyl sulfide (DAS), diallyl disulfide (DADS) and diallyl trisulfide (DAT), dithiols and ajoene. At the same time,  $\gamma$ -glutamylcysteines are converted to S-allylcysteine (SAC).<sup>9</sup> These sulphur compounds of garlic have proved to be promising antioxidants against drug induced hepatitis.<sup>11-13</sup> The liver is a vital organ. Search for new drugs for limiting hepatic injury has been of interest. The present study was aimed to explore new drug for curing acute hepatitis and delaying its progression to hepatocellular carcinoma by evaluating antioxidant and hepatoprotective properties of *Allium sativum*. Garlic is a natural component of diet in Pakistan and the Chinese exotic variety is commonly available in market. The objective of this study was to determine the hepatoprotective and antioxidative effects of ethanolic extract of *Allium sativum* var Chinese exotic on acetaminophen induced hepatotoxicity in albino rats.

## Methods

Ninety Ninety male albino rats weighing 200-250 grams from were obtained from National Institute of Health (NIH), Islamabad. Animals were housed in groups of 30 per cage for at least one week before the start of experiments. Housing conditions were thermostatically maintained at  $26 \pm 2$  °C and a light/dark cycle (lights on: 0900-2100).<sup>14</sup> The animals were fed with commercially available standard pellet diet ad libitum and were provided with tap water in clean bottles. *Allium sativum* var Chinese exotic was obtained from local market of Lahore. Ethanolic extract of *Allium sativum* var Chinese exotic was made and

standardized using facilities available at Applied Chemistry Research Centre, PCSIR labs, Lahore. The extract obtained, was filtered and the solvent (ethanol) evaporated in vacuum with a rotary evaporator. After evaporation a dark brown concentrate was obtained. This concentrate was kept at 4 °C prior to use. The crude extract was then dissolved in normal saline and then diluted to the desired concentration.<sup>15</sup>

A single intraperitoneal dose of acetaminophen 750 mg/kg<sup>16</sup> dissolved in normal saline was used to induce acute oxidative hepatic injury. Ninety male albino rats were divided into: Group A (Negative Control, n=30): was given normal saline 10ml/kg body weight intraperitoneally for 7 days. Group B (Positive Control, n=30): was given a single dose of acetaminophen 750 mg/kg<sup>16</sup> dissolved in normal saline intraperitoneally.

Group C (Experimental, n=30): was pretreated with *Allium sativum* var Chinese exotic ethanolic extract in a dose of 500 mg/kg body weight intraperitoneally<sup>17</sup> daily for 7 days before a single intraperitoneal dose of acetaminophen 750 mg/kg<sup>16</sup> dissolved in normal saline. After 24 hours of acetaminophen administration, each rat was anesthetized using ether. Three-milliliter blood was drawn by cardiac puncture and was kept in the test tube for about 15-20 minutes, and allowed to clot. After 15-20 minutes, samples were centrifuged at 5000 rpm for 15 minutes. The serum, thus obtained, was preserved in labeled polypropylene storage tubes. Serum alanine aminotransferase (ALT), aspartate aminotransferase (AST), alkaline phosphatase (ALP), total protein and albumin were determined on the same day of blood sampling. About 0.5mL of each of these samples were stored at -20 °C for determination of serum glutathione peroxidase at later stage. Data was analyzed using PASW18 (formerly SPSS). The arithmetic mean and standard deviation for quantitative variables were calculated. The statistical significance of difference amongst the three groups were determined by applying one way ANOVA followed by post hoc LSD (multiple comparison) test. The values were considered significant if the p value was less than 0.05; and, highly significant if the p value was less than 0.001.

## Results

After pretreatment with ethanolic extract of *Allium sativum* var Chinese exotic followed by acetaminophen hepatotoxicity, there was highly significant (\*p<.000) decrease in liver enzymes including serum ALT, AST and ALP in experimental group as compared to both negative and positive control groups (Table-1).

The positive control group (group B) having acetaminophen toxicity showed highly significantly ( $p=0.000$ ) raised values of serum ALT, AST and ALP as compared to the negative control group (group A) as depicted in (Table-2).

After pretreatment with ethanolic extract of *Allium sativum* Var Chinese exotic followed by acetaminophen toxicity, the experimental group C showed highly significant ( $p=0.000$ ) decrease in serum levels of ALT and AST but no significant ( $p=0.335$ ) decrease in serum level of ALP as compared to the positive control group (group B) as depicted in (Table-3).

**Table-1:** Comparison of serum ALT, AST and ALP in groups A, B and C. (One way ANOVA)

Parameters	Group A (N=30)	Group B (N=30)	Group C (N=30)	P-value
Serum ALT (U/l)	53.53±4.46	177.50±6.53	82.83±6.36	0.000*
Serum AST (U/l)	65.80±3.46	102.43±7.19	83.03±5.87	0.000*
Serum ALP (U/l)	124.30±5.81	575.90±4.69	574±2.80	0.000*

Values are presented as mean± SD \* $p<.000$  highly significant

**Table-2:** Comparison of serum ALT, AST and ALP in groups A and B. (Post hoc LSD)

Parameters	Group A (N=30)	Group B (N=30)	P-value
Serum ALT (U/l)	53.53±4.46	177.50±6.53	0.000*
Serum AST (U/l)	65.80±3.46	102.43±7.19	0.000*
Serum ALP (U/l)	124.30±5.81	575.90±4.69	0.000*

Values are presented as mean± SD \* $p<.000$ -highly significant

**Table-3:** Comparison of serum ALT, AST and ALP in groups B and C. (Post hoc LSD)

Parameters	Group B (N=30)	Group C (N=30)	P-value
Serum ALT (U/l)	177.50±6.53	87.73±3.68	0.000*
Serum AST (U/l)	102.43±7.19	61.70±4.46	0.000*
Serum ALP (U/l)	575.90±4.69	572.47±3.54	0.335**

Values are presented as mean± SD \* $p<.000$ -highly significant \*\* $p>.05$ -not significant

After pretreatment with ethanolic extract of *Allium sativum* followed by acetaminophen hepatotoxicity, there was highly significantly ( $*p<.000$ ) less decrease in serum total protein, albumin and glutathione peroxidase in experimental group as compared to both negative and positive control groups. (Table-4) The positive control group (group B) having acetaminophen toxicity showed highly significant ( $p=0.000$ ) decrease in serum total protein, albumin and glutathione peroxidase as compared to the values in negative control group (group A) as shown

in (Table-5). After pretreatment with ethanolic extract of *Allium sativum* Var Chinese exotic followed by acetaminophen toxicity, the experimental group C showed highly significant ( $p=0.000$ ) increase in serum levels of albumin, but only significant increase in total proteins ( $p=0.027$ ) and glutathione peroxidase ( $p=0.025$ ) as compared to those in the positive control group (group B) as shown in (Table-6).

**Table-4:** Comparison of serum total protein, albumin and glutathione peroxidase in groups A, B and C. (one way ANOVA).

Parameters	Group A (N=30)	Group B (N=30)	Group C (N=30)	P-value
Serum Total proteins (g/dl)	6.78±0.21	4.20±0.12	4.32±0.17	0.000*
Serum albumin (g/dl)	3.45±0.13	1.40±0.16	3.05±0.25	0.000*
Glutathione peroxidase (g/dl)	21.49±0.79	4.62±0.60	5.27±0.71	0.000*

Values are presented as mean± SD \* $p<.000$  highly significant

**Table-5:** Comparison of serum total proteins, albumin and glutathione peroxidase in groups A and B. (Post hoc LSD)

Parameters	Group A (N=30)	Group B (N=30)	P-value
Serum Total proteins (g/dl)	6.78±0.21	4.20±0.12	0.000*
Serum albumin (g/dl)	3.45±0.13	1.40±0.16	0.000*
Glutathione peroxidase (g/dl)	21.49±0.79	4.62±0.60	0.000*

Values are presented as mean± SD \* $p<.000$ -highly significant

**Table-6:** Comparison of serum total protein, albumin and glutathione peroxidase in groups B, and C. (one way ANOVA).

Parameters	Group B (N=30)	Group C (N=30)	P-value
Serum Total proteins (g/dl)	4.20±0.12	4.32±0.17	0.027***
Serum albumin (g/dl)	1.40±0.16	3.05±0.25	0.000*
Glutathione peroxidase (g/dl)	4.62±0.60	5.27±0.71	0.025***

Values are presented as mean± SD \* $p<.000$ -highly significant \*\*\* $p<.05$ -significant

## Discussion

Our study evaluated the effects of ethanolic extract of *Allium sativum* Var Chinese exotic on experimentally induced acetaminophen hepatotoxicity and noted effects on liver enzymes (ALT, AST and ALP), serum albumin, total proteins and serum glutathione peroxidase in male albino rats. This study showed that pretreatment of rats with ethanolic extract of this variety of garlic grown in Pakistan prevented the increase in liver enzymes and decrease in serum albumin, total proteins and glutathione peroxidase, due to acetaminophen toxicity. This adds to several reports on the pharmacological usefulness of garlic extracts as liver protective agents. Lee et al (June 2016) investigated the protective effect of fermented garlic extract by lactic acid bacteria (LAFGE) against acetaminophen

induced acute liver injury in rats. Their findings indicated lowered plasma ALT levels, inhibition of lipid peroxidation, glutathione and ATP depletion, and the elevation of antioxidant enzyme activities. These findings indicate that LAFGE ameliorated AAP-induced liver injury by preventing oxidative stress-mediated apoptosis, thereby establishing LAFGE as a potential supplement in the treatment of AAP-induced liver injury.<sup>18</sup> Allyl methyl disulfide (AMDS) has been identified as one of the bioactive components in fresh garlic paste and alleviates AAP-induced elevation of alanine aminotransferase (ALT), aspartate aminotransferase (AST) and lactate dehydrogenase (LDH) levels, significantly ( $p < 0.05$ ) reduces the maleic dialdehyde (MDA) level in liver tissues and restores the activities of antioxidant enzymes superoxide dismutase (SOD), glutathione peroxidase and glutathione towards normal levels.<sup>19</sup>

Ozougwu et al (2014) investigated hepatoprotective effects of *Allium sativum* methanolic extracts on paracetamol induced hepatotoxic rats. *Allium sativum* reduced ALT and total serum bilirubin in a dose dependent fashion whereas it reduced AST, ALP and LDH level in a dose independent manner. Acetaminophen hepatotoxicity leads to leakage of cellular enzymes into the plasma such as ALT, AST, ALP and LDH showing increased permeability and necrosis of hepatocytes. These significantly increased levels of serum AST and ALT are due to hepatocellular damage because these enzymes are normally located in the cytoplasm and released into the circulation after cellular damage. The mechanism of action of garlic could be through preventing the intracellular enzyme release and by membrane stabilizing effects. This is because garlic is rich in antioxidants. The reduction in ALP and LDH levels by extracts may suggest repairing of rat liver by *Allium sativum* extracts. Thus it was suggested that the active ingredients in *Allium sativum* (allyl propyl disulfide) increased the levels of glutathione to bind with the toxic metabolites of paracetamol such as N-acetyl-p-benzoquinone imine (NAPQI) and increased its rate of excretion from the body. It might also have inhibited the levels of the cytochrome P-450 enzyme system that decreased the formation of NAPQI from ingested paracetamol. These possible mechanisms of action of *Allium sativum* extracts may be through their antioxidative effects that are capable of free radical scavenging in living system.<sup>20</sup> Another study done in 2017 determined the antioxidative effects of *Allium sativum* methanolic extract against paracetamol induced liver toxicity. It was evident

that garlic extract was able to significantly raise the intracellular contents of glutathione peroxidase.<sup>21</sup>

Rashed et al (2014) investigated the effect of garlic oil (GO) alone or in combination with low dose total body gamma ( $\gamma$ )-irradiation (LDR) against paracetamol (AAP)-induced hepatotoxicity in rats. Findings showed that the combination of GO and LDR produced considerable comparable effects to either treatment alone in reducing serum elevations of ALT, AST, ALP, LDH, MDH, hepatic CYP2E1 activity and preventing the decreased hepatic glutathione content as a result of AAP toxicity. This ability of garlic to lower the raised levels of ALT, AST and to prevent decrease in levels of glutathione after AAP toxicity was in accordance with results of the present study. This remarkable synergistic protection against AAP-induced hepatotoxicity might be attributed partly to the suppressive effect of both GO constituents and LDR on lipid peroxidation by free radical scavenging properties or by restoration of glutathione content and cytochrome P450E1 enzyme in the liver.<sup>22</sup>

Shin (2014) investigated the hepatoprotective effects of aged black garlic (ABG) in rodent models of liver injury. ABG inhibited carbon tetrachloride induced elevations of ALT and AST. D-galactosamine induced hepatocellular damage was also suppressed by ABG treatment. However, ABG did not effect the elevations of ALP, which is also in accordance with the results of present study.<sup>23</sup> Sharma et al (2010) studied amelioration of lead-induced hepatotoxicity by *Allium sativum* extracts in Swiss albino mice. Oral treatment with lead nitrate induced a significant increase in the levels of hepatic AST, ALT, and ALP. Hepatic protein levels in lead-exposed mice were significantly depleted. Aqueous garlic extract and ethanolic garlic extract restored the deranged parameters significantly.<sup>24</sup> This result was in accordance with our study regarding serum parameters of ALT, AST and total proteins.

Modulatory effects of dietary inclusion of garlic (*Allium sativum*) on gentamycin-induced hepatotoxicity and oxidative stress in rats were studied by Ademiluyi et al (2013). Hepatic damage, as revealed by significant elevation of liver damage marker enzymes (AST and ALT) and reduction in plasma albumin level, were restored following consumption of diet containing garlic. Elevations of serum levels of ALT indicated necrotic lesions in the liver cells while decreased serum albumin indicated that there was an impairment in both synthetic and excretory activities of liver.<sup>25</sup> These results were in accordance with our results as ethanolic extract of garlic Chinese exotic variety prevented the decrease in plasma albumin while restoring liver damage marker enzymes to nearly normal values.

## Conclusion

Thus garlic may be considered as a useful dietary supplementary compound to patients treated with regular high doses of paracetamol such as of tuberculosis, cancer, dengue fever and arthritis. The antioxidative and hepatoprotective potential of

*Allium sativum* should be further investigated in human studies. The medical implication of this finding could be that consumption of this variety of garlic might be a useful prophylactic and therapeutic strategy against oxidative stress of toxic hepatitis in Pakistan.

## References

- Longmore M, Wilkinson I, Baldwin A, Wallin E. Oxford Handbook of Clinical Medicine. 9th ed. New York: Oxford University Press 2014.
- Friedman LS. Liver biliary tract & pancreatic disorders. In: Papadakis M.A, McPhee S.J and Rabow M.W. (editors). CURRENT Medical Diagnosis and Treatment. 57th ed. San Francisco: McGraw Hill 2018; 598-648.
- Rappaport AM. Physioanatomic considerations. In: Schiff L, Schiff ER, editors. Disease of the liver. 5th ed. Philadelphia: JB Lippincott 1982:1-57.
- Lauterburg BH, Velez ME. Glutathione deficiency in alcoholics: risk factor for paracetamol hepatotoxicity. Gut 1988; 29: 1153-7.
- Watkins PB, Kaplowitz N, Slattery JT, Colonese CR, Colucci SV, Stewart PW, et al. aminotransferase Elevations in Healthy Adults Receiving 4 Grams of Acetaminophen Daily. JAMA 2006; 296(1): 87-93.
- Bonkowsky HL, Mudge GH, Memurtry RJ. Chronic Hepatic Inflammation and Fibrosis due to low Doses of Paracetamol. Lancet 1978; 311(8072): 1016-8.
- Fontana RJ. Acute Liver Failure including Acetaminophen Overdose. Med Clin North Am. 2008; 92(4): 76194.
- Mayuren C, Reddy VV, Priya SV, Devi VA. Protective effect of Livactine against CCl4 and paracetamol induced hepatotoxicity in adult Wistar rats. N Am J Med Sci. 2010; 2(10): 4915.
- Tran GB, Dam SM, Tram Le NT. Amelioration of single clove black garlic aqueous extract on dyslipidemia and hepatitis in chronic carbon tetrachloride intoxicated swiss albino mice. Int J Hepatol. 2018; article ID 9383950.
- Mahmood T, Hussain SI, Khokhar KM, Bhatti MH, Laghari H. Comparative performance of garlic cultivars. Asian J Plant Sci 2002; 1:160-1.
- Duwairoh AM, Wirjatmadi B, Adriani M. effects of *Allium sativum* Linn. (Solo Garlic) extract in increasing superoxide dismutase (SOD) levels in *Rattus norvegicus* strain wistar exposed by E-cigarette. IJPHCS 2018; 5(4): 2289-7577.
- Singh VK, Tripathi S. hepatoprotective and antioxidative potential of *Allium sativum* in the treatment of pulmonary tuberculosis under dots. Int J Curr Res 2016; 8(3): 28202-7.
- Naji KM, Al-Shaibani ES, Alhadi FA, Al-Soudi SA, D'souza MR. Hepatoprotective and antioxidant effects of single clove garlic against CCL4-induced hepatic damage in rabbits. BMC Complement Altern Med 2017; 17:411.
- Sumioka I, Matsura T, Kasuga S, Itakura Y, Yamada K. Mechanism of Protection by S-Allylmercaptocysteine against Acetaminophen Induced liver injury in Mice. Jpn J Pharmacol 1998; 78: 199-207.
- Das Gupta A, Dhara PC, Dhundasi SA, Das KK. Effect of garlic (*Allium sativum*) on nickel II or chromium VI induced alternations of glucose homeostasis and hepatic antioxidant status under sub-chronic exposure conditions. J Basic Clin Physiol Pharmacol 2009; 20: 1-14.
- Katyare SS, Satav JG. Impaired mitochondrial oxidative energy metabolism following paracetamol - induced hepatotoxicity in the rat. Br J Pharmacol 1989; 96: 51-8.
- Drobova H, Thomson M, Al-Qattan K, Shalaby RP, Al-Amin Z, Ali M. Garlic increases antioxidant levels in diabetic and hypertensive rats determined by a modified peroxidase method. Evid Based Complement Alternat Med 2011; 2011: 703049.
- Lee HS, Lim WC, Lee SJ, Lee SH, YU H J, Lee J H, Cho H Y. Hepatoprotective effects of lactic acid-fermented garlic extract against acetaminophen-induced acute liver injury in rats. Food Sci Biotechnol 2016; 25(3): 867-73.
- Zhang Y, Zhang F, Wang K, Liu G, Yang M, Luan Y, Zhao Z. Protective effect of allyl methyl disulfide on acetaminophen-induced hepatotoxicity in mice. Chem Biol Interact 2016; 249: 71-7.
- Ozougwu JC, Eyo JE, Clarence OK, Olajotumidayo S, Kelechukwu DM. Investigation of the Antihepatotoxic Effects of *Allium sativum* Extracts Against Acetaminophen Intoxicated *Rattus norvegicus*. World J Med Sci 2014; 11(3): 397-404.
- Ozougwu JC, Eziuzor CS, Akwari DK, Ike CC. antioxidative effects of *Allium sativum* methanolic extracts against paracetamol induced liver toxicity 2017; 4(1): 24-31. 22 Rashed RR, El-Ghazaly MA, Kenawy SA. Protective effects of Garlic oil alone or combined with Low-dose Gamma Irradiation on Paracetamol-induced hepatotoxicity in rats. Eur J Biol Med Sci Res 2014; 2(3): 1-27.
- Shin JH, Lee CW, Oh SJ, Yun J, Kang MR, Han SB et al. Hepatoprotective effect of aged black garlic extract in rodents. Toxicol Res 2014; 30(1): 49-54.
- Sharma A, Sharma V, Kansal L. Amelioration of lead-induced hepatotoxicity by *Allium sativum* extracts in Swiss albino mice. Libyan J Med 2010; 5: 4621.
- Ademiluyi AO, Oboh G, Owoloye TR, Agbebi OJ. Modulatory effects of dietary inclusion of garlic (*Allium sativum*) on gentamycin-induced hepatotoxicity and oxidative stress in rats. Asian Pac J Trop Biomed 2013; 3(6): 470-75.