

Original Article

ASSESSMENT OF CASES OF CONGENITAL CATARACT IN PATIENTS ADMITTED IN HOSPITALS OF LAHORE

Muhammad Naeem, Najam Ud Din, Malik Shahid Shaukat, Khaleeq Ahmad Qureshi, Irfan Yameen and Sahar Fatima

Objective: To observe the different patterns of congenital cataract in children and to determine the age at which maximum cases are reported.

Material and Methods: Descriptive cross sectional .Study was conducted on patients of congenital cataract in different hospitals of Lahore and Study duration was 03 months. All participants were selected by non probability purposive sampling technique. Parents of patients of congenital cataract were interviewed after taking verbal consent using questionnaire method. Data was analyzed using SPSS.

Results: In this study 30 cases of congenital cataract were examined age ranging from 0-8 years in the four hospitals of Lahore. Majority (70%) of the cases of congenital cataract are reported up to age 4 years. Out of 30 cases of congenital cataract 56% are male and 43% are female. Majority (83.3%) of cases are reported with bilateral congenital Cataract. Nystagmus and squint as associated symptoms are present in majority (63.3%) cases. Hereditary trait regarding congenital cataract is observed in 46.7% of cases. Maternal rubella infection as a cause of congenital cataract is reported in majority (73.3%) of cases. Congenital cataract in associated with metabolic disorders is reported in 26.7% of cases. Majority of patients (73.3%) are having lamellar pattern.

Conclusion: Most patients are reported up to the age of 4 years. Males are predominantly affected. Maternal rubella is found to be commonest cause. Most of cases are with bilateral presentation. Majority of cases have lamellar pattern.

Introduction

Cataract is any light scattering opacity of lens present at birth. 2/3rd are bilateral, most of them are due to autosomal dominant mutation, unilateral cataracts are sporadic, only 10% have a definite cause. Infantile prevalence in U.S.A showed a rate of 3-4 visually significant cataracts per 10000 live births.¹ Common causes are hypoglycemia, trisomy, myotonic dystrophy, infectious diseases {TORCH}, prematurity. About fifteen genes involved in cataract formation have been identified and the inheritance is most often autosomal dominant, although it can be x-linked or autosomal recessive.² High summer heat upto 5 degree increase in apparent temperature may cause congenital cataract with 51% increase in the risk of congenital cataract.³ Cataracts are felt to be visually significant. If the treatment is delayed it can lead to amblyopia and visual loss can occur in later stages .Surgical intervention is the only option, the timing of surgery is critical for visual development, most investigators recommend surgery within the first two months of life.⁴ There has been evidence to suggest that before one month of age the risk of aphakic glaucoma is increased.⁵ Intraocular lens implantation in children is considered to be safe and

acceptable in children.⁶

Material and Methods

It was a descriptive cross sectional study conducted in Services Hospital, Lahore, Jinnah Hospital, Lahore, Mayo Hospital, Lahore, Rehmat Begum Trust Hospital, Lahore. The study duration was three months. The study subjects were children up to 8 years of age. Patients with acquired cataract were excluded from the study. All the parents of the subjects were explained the purpose and process and benefits of the study. Confidentiality was ensured. A check list and a questionnaire was developed. Cataract was diagnosed by screening eye examination including.

- 1) Red reflex test by direct ophthalmoscope.
- 2) Slit lamp examination.

A Sample of 30 patients was selected by non probability purposive sampling technique. Data was analyzed by SPSS computer software.

Results

In our study we observed that 70% of the reported cases of congenital cataract were up to age 4 years. Out of 30 cases of congenital cataract, 56% were

male and 43% were female (**table-1**). It was observed that majority (83.3%) reported with bilateral congenital Cataract (**table-3**). Nystagmus and squint as associated symptoms was present in majority (63.3%) cases. Hereditary trait regarding congenital cataract was observed in 46.7% of cases. Maternal rubella infection as a cause of congenital

cataract was reported in majority (73.3%) of cases. Congenital cataract in associated with metabolic disorders was reported in 26.7% of cases. On slit lamp examination of cases of congenital cataract. It was observed that majority of patients (73.3%) were having lamellar pattern (**table-4**).

Table-1: Slit lamp pattern presentation of patient gender of patient cross tabulation count.

Gender of Patients			Presentation of Patient		Total
			One eye	Both eyes	
Male	Slit Lamp Pattern	Lamellar	02	11	13
		Posterior Capsular	02	02	04
		Total	04	13	17
Female	Slit lamp Pattern	Lamellar	0	09	09
		Posterior Capsular	01	03	04
		Total	01	12	13

Table-2: Age of presentation in years(grouped) associated symptoms of patient cross tabulation count.

		Associated symptoms of patients		
		Present	Absebt	Total
Age of presentation in years	4 or less	13	08	21
	5 or more	06	03	09
Total		19	11	30

Table-3: Presentation of patient mode of delivery cross tabulation count.

		Mode of delivery		Total
		Normal	C-section	
Presentation of patients	One eye	05	0	05
	Both eyes	18	07	25
Total		23	07	30

Table-4: Slit lamp pattern presentation of patient age of presentation in years(grouped) cross tabulation.

Age of presentation in years (grouped)			Presentation of Patient		Total
			One eye	Both eyes	
4 or less	Slit Lamp Pattern	Lamellar	02	13	15
		Posterior Capsular	02	04	06
		Total	04	17	21
5 or more	Slit lamp Pattern	Lamellar	0	07	07
		Posterior Capsular	01	01	02
		Total	01	08	09
Total	Slit lamp Pattern	Lamellar	02	20	22
		Posterior Capsular	03	05	08
Total			05	25	30

compilation and analysis.




Fig-1: Age of presentation.

Discussion

Congenital cataract is responsible for nearly 10% of all vision loss in children world wide.¹⁶ Their early detection and treatment holds a vital position in the prevention of permanent visual loss. In this study, 30 cases of congenital cataract were examined age ranging from 0-8 years in the four hospitals of Lahore. In our study we observed that 70% of the reported cases of congenital cataract were up to age 4 years.

Out of 30 cases of congenital cataract 56% were male and 43% were female (**table-1**), this is in accordance with the research carried out in West Bank and Gaza Strip.¹⁷ It was observed that majority (83.3%) were reported with bilateral congenital Cataract (**table-3**). It is in accordance with the result of a research carried out in United Kingdom.¹⁸

Nystagmus and squint as associated symptoms was present in majority (63.3%) cases and is in accordance with research carried out in United States of America that these are most common ocular anomalies associated with congenial cataract.¹⁹ Hereditary trait regarding congenial cataract was observed in 46.7% of cases. According to research carried out in china upto 25% of cases of congenial cataract are believed to be inherited.²⁰

Maternal rubella infection as a cause of congenial cataract is reported in majority (73.3%) of cases. This result is in contraindication with research carried out in South India.²¹ Higher incidence of maternal rubella infection in Pakistan is due to unhygienic delivery practices and lack of awareness about rubella vaccination. Congenial cataract in associated with metabolic disorders is reported in 26.7% of cases. It is in accordance with the research carried out in U.S.A which shows that only few cases of congenial cataract are associated with metabolic disorders.²²

After slit lamp examination of cases of congenial cataract it was observed that majority of patients (73.3%) were having lamellar pattern.

Conclusions

Most patients were reported up to the age of 4 years. Males were predominantly affected. Maternal rubella was found to be commonest cause of congenial Cataract due to septic birth practices. Majority of cases of congenial cataract were with bilateral presentation. Majority of cases had lamellar pattern on slit lamp examination.

*Department of Community Medicine
Services Institute of Medical Sciences Lahore*

References

- Holmes JM, Leske DA, Burke JP and Hodge DO. Birth prevalence of visually significant infantile cataract in a defined U.S. population. *Ophthalmic Epide- miology* 2003 Apr; 10:67-74
- Ashwin R M, Molecular Genetic Basis of Inherited Cataract and Associated Phenotypes. *Survey of Ophthalmology* 2004 May-June
- van Zutphen ,AR,S Lin,BA Fletcher and S-A Hwang. population based case-control study of extreme summer temperature and birth defects. *Survey of ophthalmology* 2012
- Pandey SK, Wilson ME, Trivedi RH, et al. Pediatric cataract surgery and intraocular lens implantation: current techniques, complications and management. *Int Ophthalmol Clin.* 2001 Summer;41(3):175-96.
- Vishwanath M, Cheong-Leen R, Taylor D, et al. Is early surgery for congenital cataract a risk factor for glaucoma? *Br J Ophthalmol* 2004;88:905-910.
- Infant Aphakia Treatment Study Group, Lambert SR, Buckley EG, Drews-Botsch C, Dubois L, Hartmann EE, Lynn MJ, Plager DA, Wilson ME. A randomized clinical trial comparing contact lens with intraocular lens correction of monocular aphakia during infancy: grating acuity and adverse events at age 1 year. *Arch Ophthalmol.* 2010. Jul;128:810-8.

7. Ashwin Reddy M, et al. Molecular Genetic Basis of Inherited Cataract and Associated Phenotypes. Survey of Ophthalmology 2004 May-June
8. Pandey SK, Wilson ME, Trivedi RH, et al. Pediatric cataract surgery and intraocular lens implantation: current techniques, complications and management. Int Ophthalmol Clin. 2001 Summer;41(3):175-96.
9. Tesser RA, Hess DB, Buckley EG. Pediatric cataracts and lens
10. Anomalies. In: Harley's Pediatric Ophthalmology, 5th ed, Nelson LB, Olitsky SE (Eds), Lippincott Williams & Wilkins, Philadelphia 2005. p.255.
11. Wolff SM. The ocular manifestations of congenital rubella. Trans Am Ophthalmol Soc 2005; 70:577.
12. World Health Organization Office of Information. Blindness and visual disability: Part 2: Major causes worldwide.. WHO fact sheet 143. Geneva: WHO, 1997
13. Fraser FC, Sajoo A. (1995) Teratogenic potential of corticosteroids in humans. Teratology;51:45-6.
14. Acta Anat (Basel). 1990;137(2):132-6.
15. Snyder C. Aurelius Cornelius Celsus On Cataracts. Arch Ophthalmol. 1964 Jan;71:144146. [PubMed]
16. Santana, A; Waiswo, M (2011 Mar-Apr). "The genetic and molecular basis of congenital cataract." Arquivos brasileiros de oftalmologia 74 (2): 13642. PMID 21779674.
17. Lambert SR. congenital cataract predominancy in males . In: Pediatric Ophthalmology, Neuro-Ophthalmology, Genetics, Lorenz B, Moore AT (Eds), Springer, New York 2006. p.81.
18. Jugnoo S. Rahi, Department of Epidemiology and Public Health, Institute of Child Health, 30 Guilford Street, London WC1N 1EH, UK. j.rahi@ich.ucl.ac.uk. Submitted for publication October 20, 1999; revised February 4, 2000; accepted February 18, 2000.
19. Rahi JS, congenital and infantile cataract, London, united kingdom. invest Ophthalmol Vis Sci. 2000 july;41(8):2108-14
20. Hejtmancik JF. Congenital cataracts and their molecular genetics. Semin Cell Dev Biol. 2008;19:13449. [PMC free article][PubMed]
21. Vijayalakshmi P, Rajasundari TA, Prasad NM, Prakash SK, Narendran K, Ravindran M, Muthukkaruppan VR, Lalitha P, Brown DW. Aravind Eye Hospital, 1, Annanagar, Madurai 625 020, India. P.vijayalakshmi@aravind.org Br J Ophthalmol. 2007 Nov;91(11):1467-70.
22. The American Association for Pediatric Ophthalmology and Strabismus Annual Meeting, April 2009, San Francisco, CA. Sasapin G. Pralapakorn, MD, MPH.